

Table 1. Arkansas lime recommendations for lawns.

Soil Texture	Soil Water pH Level Interpretation					
	Below Optimum			Medium	Optimum	High
Soil Test Ca	< 5.0	5.0 - 5.4	5.5 - 5.7	5.8 - 6.2	6.3 - 6.9	> 6.9
ppm	----- lbs CaCO ₃ lime/1,000 ft ² -----					
< 500	80 [†]	57	46	0	0	0
500 - 1500	115	92	69	0	0	0
1501 - 2250	138	115	92	0	0	0
> 2250	138	138	115	0	0	0

[†]Lime rates are for CaCO₃ (Ag lime) sources that contain a 90 percent calcium (Ca) carbonate equivalent. Do not apply more than 50 pounds CaCO₃ lime/1,000 ft² in any one application. Water-in lime after application to avoid foliar burn. If magnesium (Mg) is recommended by a soil test, then use dolomitic lime (MgCO₃ • CaCO₃) or a lime with 10 percent or more magnesium instead of calcitic lime (CaCO₃) (< 2% Mg). Please contact your local extension specialist to determine how much lime is needed if using another source of lime such as hydrated lime. Collect soil samples every one to two years to monitor soil pH.

Each turfgrass species prefers a slightly different pH for optimum growth. Bermudagrass, tall fescue and zoysiagrass prefer a soil pH in the range of 5.8 to 7.0. St. Augustinegrass prefers more alkaline conditions with an optimum pH of 6.5 to 7.5. Centipedegrass is more tolerant of acid soils (pH = 5.0 to 6.0) compared to other species. Lime should not be applied to centipedegrass lawns unless soil pH is < 5.0, whereas other turfgrass species will benefit from lime if soil pH drops below 5.8.

Turfgrass health will improve with lime applications if your soil pH is less than 5.8 and your lawn species is not centipedegrass. Your soil pH and the amount of calcium already on your soil's cation exchange sites provide the information necessary to determine how much lime you should apply to increase your soil pH (Table 1). When you receive the results of your soil test, the soil test report will indicate whether or not your lawn needs liming and how much lime to apply. Lime is best applied in the spring or fall, but regardless of application timing, it may take several months before the lime can effectively raise the pH of your soil. It is important not to apply more than 50 pounds/1,000 ft² in any one application when applying lime. Additionally, irrigating the lawn after liming will help reduce the risk of any foliar burn from the lime.

If it is recommended that you apply < 50 pounds lime/1,000 ft², apply this in the spring or fall as a single application. If between 50 and 100 pounds lime/1,000 ft² is recommended, apply half the recommended rate in the spring and the other half in the fall. If more than 100 pounds lime/1,000 ft² is recommended, apply 50 pounds in the spring and 50 pounds in the fall. Continue to lime and retest your soil annually until the pH is optimum for turf growth.

There are many lime choices available to the homeowner. They contain different quantities of

calcium and magnesium. Additionally, these products have different particle (mesh sieve) sizes that influence how quickly these products affect soil pH. The quality of lime is determined by various calculations, taking into consideration the fineness and nutrient content. Each product should include information about calcium and magnesium content as well as fineness (Table 2). State lime regulatory laws vary, so each state measures lime quality differently. Regardless of the state where the lime is manufactured, a good rule of thumb is to choose a product that has a relative neutralizing value (RNV) or effective calcium carbonate equivalent (ECCE) > 80 percent in order to ensure that the lime is of good quality.

Table 2. Information commonly included on a lime label.

Minimum guaranteed screen analysis	
Screen size	% Passing
8 mesh	100
10 mesh	100
20 mesh	95
40 mesh	90
50 mesh	85
60 mesh	80
100 mesh	70

Minimum guaranteed chemical analysis (dry)	
Calcium (Ca)	21.6%
Magnesium (Mg)	10.0%
Calcium carbonate equivalent (CCE)	95.0%
Relative neutralizing value (RNV)	92.0%
Effective calcium carbonate equivalent (ECCE)	86.0%

Liming materials with a fine particle size such as pulverized limestone may be difficult to distribute accurately with a rotary spreader but can be more easily distributed with a drop spreader (Figures 2 and 3). Lime can be pelletized so that many small particles are bonded together, which allows the product to be more easily applied (Figure 4). Pelletized liming materials can be distributed with a rotary spreader or a drop spreader.



Figure 2. Comparison of pelletized lime (left) and pulverized limestone (right).

Liming Terminology

1. **CCE:** Calcium carbonate equivalent. Relative purity of the liming material using pure calcium carbonate (CaCO_3) as the standard.
2. **ECCE:** Effective calcium carbonate equivalent. An expression of lime effectiveness based on the combined effect of chemical purity (CCE) and fineness (sieve size). This calculation method differs between states.
3. **Lime:** A soil amendment that contains calcium carbonate (CaCO_3), magnesium carbonate (MgCO_3) and other materials. Lime is used to neutralize soil acidity and provide calcium and magnesium, macronutrients required for plant growth.
4. **pH:** Negative logarithm of the hydrogen ion concentration. It is a measurement of acidity or alkalinity.
5. **RNV:** Relative neutralizing value. An expression of lime effectiveness based on the combined effect of chemical purity (CCE) and fineness (sieve size). This calculation method differs between states.



Figure 3. Pulverized limestone has a very fine texture and is more easily applied with a drop spreader.



Figure 4. Pelletized lime can be applied with either a drop or rotary spreader.

Additional fact sheets available at <http://publications.uaex.edu/>.

For more information about turfgrass, visit <http://turf.uark.edu/>.

References

1. Carrow, R.N., D.V. Waddington and P.E. Rieke. 2001. *Turfgrass Soil Fertility and Chemical Problems: Assessment and Management*. Ann Arbor Press, Chelsea, MI.
2. Espinoza, L., M. Mozaffari and N.A. Slaton. 2007. University of Arkansas lime and fertilizer recommendations. University of Arkansas Cooperative Extension Service.
3. Johnson, B.J., and R.N. Carrow. 1992. Influence of soil pH and fertility programs on centipedegrass. *Agron. J.* 84:21-26.
4. Slaton, N.A. 2007. Associate professor of Soil Fertility and director of Soil Testing, University of Arkansas, Fayetteville. Personal communication.

Printed by University of Arkansas Cooperative Extension Service Printing Services.

DR. AARON PATTON is an assistant professor - turfgrass specialist with the University of Arkansas Division of Agriculture, Cooperative Extension Service, Fayetteville.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director, Cooperative Extension Service, University of Arkansas. The Arkansas Cooperative Extension Service offers its programs to all eligible persons regardless of race, color, national origin, religion, gender, age, disability, marital or veteran status, or any other legally protected status, and is an Affirmative Action/Equal Opportunity Employer.

FSA6134-PD-8-07RV